


Effective *SOD1* targeting with vMiX™, an innovative AAV-based RNA interference platform

L. Reilly,¹ Z. Walker,¹ D. Ojutalayo,¹ C. Andreassi,¹ N. Sheikh,¹ E. Dellow,¹ N.A. Mohammadi Nafchi,² O. Brock,² A. Abhilash,¹ L. Ireland,¹ R. Elsayed,¹ A. Ule,¹ P.M. Coan,¹ C.J. Miranda,¹ J. Isaac,¹ C.E. Shaw,^{1,2} D.Y. Lee,¹ Y.B. Lee,^{1,2} R. Joubert¹

¹AviadoBio Ltd, London, UK; ²Maurice Wohl Clinical Neuroscience Institute, King's College London, London, UK

AVIADOBIO



vMiX™


P0280

OBJECTIVE

To design, screen, and select effective microRNA (miRNA) candidates using the vMiX™ platform for efficient silencing of *SOD1*, a causative gene in amyotrophic lateral sclerosis (ALS).

Lead candidate selection screening of *SOD1* miRNA candidates

Figure 4: *In vitro* screening of candidate miRNAs targeting *SOD1*




- A)** miRNAs targeting *SOD1* in human and additional species were designed and cloned into the vMiX™ vector. HEK293 cells were transfected, and RNA harvested as before. All candidates successfully induced knockdown of *SOD1*, varying from 38% to 65% reduction in expression compared to a non-targeting miRNA control (Figure 4A; one-way ANOVA with Dunn's multiple comparisons, **p<0.01).
- B)** HEK293 cells were transfected and cultured for 72 hours before harvesting and protein extraction. All candidates induced knockdown of *SOD1*, from 21% to 35% reduction in protein levels compared to a non-targeting miRNA control (Figure 4B).

INTRODUCTION

- ALS is a severe neurodegenerative disease characterized by progressive motor neuron loss, leading to muscle atrophy, paralysis, and respiratory failure. *SOD1*, the first gene linked to ALS, accounts for 12–20% of familial ALS and 1–3% of sporadic ALS cases. Toxic gain-of-function *SOD1* mutations cause protein misfolding and aggregation.
- Recent approval of Qalsody® (tofersen) validates gene silencing as a therapeutic treatment approach for *SOD1*-ALS. However, the need for frequent, chronic antisense oligonucleotide injections creates significant treatment burden for patients and caregivers, and also non-trivial associated healthcare resource utilization. Consequently, there is interest in long-term, potentially one-time treatments that could meaningfully reduce treatment burden over time.
- Using vMiX™, a novel adeno-associated virus (AAV)-based RNA interference platform utilizing miRNA for gene silencing (see Poster P0944), we aimed to design, screen, and select effective miRNA candidates for efficient *SOD1* knockdown, evaluating their performance *in vitro* and *in vivo*.


In vitro screening of *SOD1* miRNA candidates

Figure 1: *In vitro* development of miRNA targeting *SOD1* using a luciferase assay




- miRNAs targeting across the coding sequence of *SOD1* were designed and cloned into a reporter plasmid and co-transfected in HEK293 cells.
- 24 hours post-transfection, a luminescence assay was performed.
- Certain miRNA sequences reduced detected levels of luminescence by up to 90% (Figure 1; SD9; Kruskal-Wallis with Dunn's multiple comparisons, *p<0.05, **p<0.01).

Figure 2: *In vitro* analysis of *SOD1* knockdown following transfection of the vMiX™ vector




- Eight miRNAs which most effectively reduced luminescence were cloned into the vMiX™ vector to assess knockdown of endogenous *SOD1*.
- HEK293 cells were transfected with vMiX™ vectors and cultured for 48 hours before harvesting and RNA extraction.
- All candidate miRNA successfully induced knockdown of *SOD1*, varying from 65% to 90% reduction in expression compared to a non-targeting miRNA control (Figure 2; SD9; Kruskal-Wallis with Dunn's multiple comparisons, **p<0.01).

Figure 3: *In vitro* analysis of *SOD1* knockdown following transduction of the vMiX™ vector




- AAV9 vectors were produced for the three miRNAs which most effectively reduced *SOD1* expression.
- HEK293 cells were transduced with vMiX™ vectors and cultured for 48 hours before harvesting and RNA extraction.
- Candidate miRNAs reduced expression of *SOD1* by up to 57% (Figure 3; SD9; Kruskal-Wallis with Dunn's multiple comparisons, *p<0.05, **p<0.01).

Figure 5: *In vitro* transduction with candidate miRNA guides targeting *SOD1*




- AAV9 vectors were produced for five miRNAs which effectively reduced *SOD1* expression. HeLa cells were transduced with vMiX™ vectors and cultured for 72 hours before harvesting and RNA extraction.
- Candidate miRNAs reduced expression of *SOD1* by up to 70% (SD46) or 81% (SD9; Figure 5; two-way ANOVA with Dunnett's multiple comparisons, **p<0.01; ***p<0.001; ****p<0.0001).

Figure 6: *In vivo* confirmation of *SOD1* knockdown with vMiX™



- A)** Neonatal mice expressing the human mutated G93A *SOD1* received intracerebroventricular injections of a vMiX™ AAV9-miR-*SOD1*. After six weeks, analysis of cortical samples demonstrated an efficient knockdown of *SOD1* mRNA (Figure 6A; one-way ANOVA with Dunnett's multiple comparisons, ****p<0.0001).
- B)** Small RNA sequencing miRNA processing analysis revealed candidate guides produced a similar average amount of guide species and low amounts of passenger. SD46 displayed the most accurate guide processing at the expected 21nt length (data not shown).



- C)** Analysis of cortical samples demonstrated an efficient knockdown of *SOD1* mRNA across a broad range of vector genome/dg quantifications. SD9 and SD46 display correlations between the amount of guide expressed and vg/cell quantifications (data not shown).

CONCLUSIONS

- We have developed and demonstrated that the novel RNA interference platform vMiX™ has the capacity to efficiently silence a gene associated with both familial and sporadic ALS.
- This platform shows broad adaptability for an AAV-based RNA interference approach to target a range of diseases.